

# AMICOS KPM

AMICOS KPM IS A BROAD SPECTRUM, CONTACT FOLIAR FUNGICIDE, FOR CONTROL OF A VARIETY OF DISEASES ON AGRICULTURAL CROPS, TERRESTRIAL AND ORNAMENTAL PLANTS AND GREENHOUSE CROPS AND ORNAMENTALS

**ACTIVE INGREDIENT:**

Sodium chloride í .. 1.0%

**OTHER INGREDIENTS:** í 99.0%

Potassium bicarbonate

Kaolin

**TOTAL:** í .. 100.0%

Net content: X Pounds

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

**FIRST AID**

**IF IN EYES:** Flush with plenty of water. Seek medical attention if irritation persists.

**IF ON SKIN:** Wash with soap and water.

Take the product's label or container with you when seeking medical attention.

**PRECAUTIONARY STATEMENTS**

Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Persons handling concentrate are recommended to wear:

• Protective eyewear

• Chemical resistant gloves made of neoprene, nitrile or natural rubber

**User Safety Recommendations**

User should wash its hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if the product gets inside. Wash thoroughly and put on clean clothing. Remove PPE after handling this product. Wash the outside of the gloves before removing.

**DIRECTIONS FOR USE**

Mode of Action: AMICOS KPM is a contact fungicide, so direct contact with the fungus organism is required for control

AMICOS KPM must be applied using a sufficient volume of water to insure complete coverage of all stems and foliage. Final spray solution should not be below 7.0 pH.

**SPRAY APPLICATION METHOD:** AMICOS KPM may be applied using conventional ground or hand held application equipment. Use quantities of water sufficient to provide thorough coverage of infested plants. Do not apply this product through any type of irrigation system. For better results, use a properly calibrated sprayer and check the calibration frequently.

**Always follow these directions:**

Spray directly onto pest(s) or pathogen(s).

Apply before noticeable foliar damage occurs.

Thorough spray coverage is essential for desired pest control.

**Compatibility**

AMICOS KPM has not been tested for use in a tank mix with many commonly used insecticides, fungicides, or spray tank adjuvants. We recommend testing all tank mix combinations for physical compatibility and plant safety. Consult your dealer or distributor for specific recommendations. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not exceed label dosage rate. As with any pesticide, the addition of spray adjuvants and other pesticides to the spray tank may cause phytotoxicity to the foliage and/or fruit of the susceptible crops and plants.

### Mixing Instructions

AMICOS KPM must be diluted with water before applying. Pour required amount into spray tank partially filled with water. Add AMICOS KPM slowly to the tank while hydraulic or mechanical is running and continue filling with water. Add spreader/sticker and/or other products last. Do not allow mixture to stand. Use mixture within 12 hours as AMICOS KPM may lose its activity if left in the tank.

### APPLICATION RATE TABLE

Spray volume per acre of treatment can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. For the control of powdery mildew, use AMICOS KPM as indicated in the following crops. Use the higher rate when disease pressure increases or a severe infection has occurred. Repeat application at 10-14 day intervals or as needed. Increase frequency of application to every 5-7 days during periods of weather favorable to fungal growth. **DO NOT EXCEED A MIX RATE OF 5 POUNDS OF THIS PRODUCT PER 100 GALLONS OF WATER.** Do not store unused spray solution. Dispose of unused spray solution on site. Spray solution should be applied within 12 hours of preparation.

CROPS	RATE of AMICOS KPM
<b>Root and tuber vegetables</b> , such as potato, beet (garden sugar), carrot	2.5 to 3.0 lbs/A
<b>Bulb vegetables</b> , such as onions (dry bulb, green), garlic, leeks, shallots.	
<b>Leafy vegetables</b> , such as endive, head and leaf lettuce, radicchio, spinach, Swiss chard.	
<b>Cole crops</b> , such as broccoli, Brussels sprouts, cabbage, cauliflower.	
<b>Citrus</b>	
<b>Legume vegetables</b> (succulent or dried), such as bean (snap, green), pea (edible-pod, field, green, snow).	
<b>Fruiting vegetables</b> , such as eggplant, pepper (bell, chilli, pimento), tomatillo, tomato.	
<b>Pome fruits</b> , such as apple, pear, quince.	
<b>Stone fruits</b> , such as apricot, cherry, nectarine, peach, plum, prune.	
<b>Small fruits and berries</b> , such as blackberry, blueberry, currant, loganberry, raspberry, strawberry.	
<b>Subtropical fruits</b> , such as banana, date, mango, papaya, pineapple.	
<b>Cereal grains</b> , such as barley, wheat.	
<b>Herbs and spices</b> , such as basil, chives, coriander (cilantro), dill, oregano, parsley, thyme.	
<b>Nut Crops</b> , such as Almond, Walnut, Pecan, Pistachio, hazelnut.	
<b>Avocado</b>	
<b>Additional crops</b> , mint (peppermint and spearmint), tobacco.	1.0-3.0 lbs/A or 1-3 lb/100 gal. water/A for hand held applications
<b>Ornamentals-field</b> , such as bare root, container, bedding and flowering stock, field grown cut flowers, vegetable transplants, nursery and landscape, potted flowering, shade and flowering trees, woody ornamentals	
<b>Roses</b>	2.5-3.0 lbs/A or 1-3 lb/100 gal. water/A for hand held applications
<b>Turf</b> (ornamental lawns, golf courses and sod farms)	
<b>Cucurbits</b> , such as cucumber, cantaloupe, casaba, honeydew, melon, pumpkin, summer squash (crookneck, zucchini), winter squash (butternut, acorn), watermelon.	2.5-5 lbs/A
<b>Hops</b>	
<b>Grapes</b> (wine, fresh market, raisin).	

**Note to User:** Unintentional consequences such crop injury may result due to environmental or growing conditions, manner of use or application. Because plant tolerance of pesticides varies as conditions vary, treat a few plants under conditions unexpected to be encountered and observe for plant damage prior to full-scale application to large number of plants.

**STORAGE AND DISPOSAL**

Pesticide Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fullerø earth, etc. Dispose of with trash.

**WARRANTY AND CONDITIONS OF SALE**

SELLER OFFERS AND THE BUYER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED TERMS AND CONDITIONS OF SALE AND WARRANTY

The directions for the use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all the variables and to eliminate risks associated with its use. Injury or damage may result because of conditions that are beyond the control of the SELLER.

SELLER warrants that this product conforms to the chemical description on this label and is believed to be reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under conditions of normal use.

SELLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

To the fullest extent permitted by law, the SELLER shall not be liable for consequential, special or indirect damages resulting from the use or handling of the product. Any variation or exemption from this warranty must be in writing and signed by authorized representative or the seller.

This product has not been registered by the U.S. Environmental Protection Agency (EPA). SEIPASA SA represents that this product qualifies for exemption from registration under section 25(b) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Manufactured by:

SEIPASA SA

C/Almudevar 22240 Tardienta (Spain) tel. +34974253433 fax +34974253298

[www.seipasa.com](http://www.seipasa.com)

Made in Spain