



HOPS

For increased Early Season Vigor, Improved Growth & Yield

Myconate activates mycorrhiza, enhancing early season root growth and colonization leading to improved crop vigor and thus profitability of hop production

Use Recommendations: Hops

- Application Timing:** Apply Myconate at planting or within 1 week of transplanting. This application stimulates the Mycorrhiza in the root zone to germinate and colonize the roots for improved early season vigor and season long benefits.
- Banded Application:** Apply 2.75 oz Myconate AS in a band or to hills with-in one week before or after transplanting. Follow application with sufficient water to move product into the root zone
- In Furrow Application:** Apply 2.75 oz Myconate AS at planting in seeding furrow within 1 inch of transplant placement.
- Transplant Drench:** Apply 2.75 oz of Myconate AS diluted in enough solution to drench the roots on the number of transplants to be planted on each acre.
Transplant within a week of application
- Overhead Irrigation:** Apply 2.75 oz of Myconate AS per acre though overhead irrigation system within 1 week of planting
- Drip Line Application:** Apply 2.75 oz of Myconate AS per treated acre within 1 week after planting or transplanting for best results
- For Best Performance:** Apply Myconate with in or very near the root zone early in growing season
- Other Considerations:** Myconate is compatible with most pesticides, and fertilizers
- Regulatory:** Myconate is water based soil additive and is not a pesticide or fertilizer
Read and follow label directions before using this product



Sustainable Solutions for Agriculture

111 S Court Str. Ste. 206
Visalia, CA 93291
www.sym-agro.com