



Vegetables *High Density*

For Increased Crop Yield and Early Vigor:

Myconate activates native mycorrhiza, enhancing early season root growth and colonization leading to improved crop vigor and thus profitability of vegetable production

Use Recommendations: Onions, Leeks, Lettuce, Celery, Carrots

- Application Timing:** Apply Myconate to seed, at planting or with in 1 week of transplanting, to stimulate the Mycorrhiza in the root zone to germinate and colonize the roots for improved early season vigor and season long benefits.
- Seed Treatment:** Apply 1.25 oz **Myconate HB** per acre worth of seed at or before planting.
- Banded Application:** Apply 2.5 oz **Myconate AS** in a band with-in one week before or after seeding or transplanting. Follow application with sufficient water to move product into the root zone
- In Furrow Application:** Apply 2.5 oz **Myconate AS** at planting in seeding furrow with in 1 inch of seed
- Transplant Drench:** Apply 2.5 oz of **Myconate AS** diluted in enough solution to drench or dip on the number of transplants to be planted on each acre. For transplant flats water with ¼ - ½ inch of water to incorporate into root zone after application. Transplant with in a week of application
- Overhead Irrigation:** Apply 2.75 oz of **Myconate AS** per acre though overhead irrigation system with in 1 week of planting
- Drip Line Application:** Apply 2.5 oz of **Myconate AS** per treated acre with in 1 week after planting or transplanting for best results
- For Best Performance:** Apply Myconate with in or very near the root zone early in growing season
- Other Considerations:** Myconate is compatible with most pesticides, and fertilizers
- Regulatory:** Myconate is water based soil additive and is not a pesticide or fertilizer