



ProBlad[®] Verde

Attack fungal diseases with an entirely new mode of action

Labeled Crops

- Grapes
- Almonds
- Strawberries
- Tomatoes
- Stone Fruit*
- Pome Fruit*
- Fruiting vegetables*
- Cucurbits*
- Hops*

*Not registered for use in California

Labeled Diseases

- Powdery mildew
- Botrytis
- Anthracnose
- Brown rot blossom blight

Key Features

- Fast acting, stops disease in 4-8 hours
- Controls all disease life stages, including spores
- New multi-site mode of action
- Translaminar activity
- Curative/reach back activity for up to 7 days
- Provides 10-14 day spray interval
- Breaks disease resistance
- Minimal toxicity to adult bees, bee larvae, and beneficial predators

ProBlad[®] Verde fungicide introduces a completely new mode of action to disease management, providing decisive, multi-process disruption in a way no other fungicide can claim. Formulated with Blad polypeptide technology, it is so new that a new FRAC Group code was created for ProBlad fungicide's active ingredient.

In this way, ProBlad fungicide separates itself from all other fungicides and has no known cross-resistance to any fungicide class, making it the optimum resistance management partner.

ProBlad fungicide works on contact by binding to the fungal cell wall and deforming the chitin structures within, ultimately tearing the fungal cell apart. This action then starts a chain of events that disrupt nutrient absorption, end cellular reproduction and destroy the fungal cell. All this occurs within eight hours of contact. The result is quick, reliable disease control that meets or exceeds established standards.

Bottom line, ProBlad fungicide is driven by a novel, multiple-site mode of action, has proven results and protects crops today with a new mode of action that extends the useful life of existing chemistries.

Application Guidelines

Preharvest Interval: One day

Re-Entry Interval: Four hours

Maximum Amount of ProBlad[®] Fungicide Per Season: No more than five foliar applications per crop season or more than two sequential applications

Label

Botrytis Rates: 24.4–36.6 fl. oz./A

Powdery Mildew Rates: 20.5–24.4 fl. oz./A

Brown Rot Blossom Blight Rates: 20.5-24.4 fl. oz./A

Active Ingredient: Banda de Lupinus albus doce (BLAD)

Mode of Action: Inhibition of cell wall synthesis and nutrient uptake by binding to chitin structures and rupturing fungal cell

FRAC Group: BM01

SymAgro
www.sym-agro.com

1 Day PHI
4 Hour REI
MRL exempt



541-607-5097
Info@Sym-Agro.com