

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE IN ALL STATES**CINNEX FUNGICIDE INSECTICIDE**

EPA Reg. No.: 87946-1

FOR THE PREVENTION AND CONTROL OF MITES AND BACTERIAL DISEASES

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Pest	Rate (fl. oz/100 gallons)	Dilution (Gal/Acre)	Timing/Application
ACARINA including but not limited to: mites* including, rust mite, red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, Pacific spider mite and two-spotted spider mite. *Not intended for use on public health pests.	32-64	50 – 300 by ground 7-20 by air	Apply in sufficient water to provide full coverage Begin applications at the first signs of mites, continue applications on a 7 day spray interval
Including but not limited to, Almond Leaf Scorch (<i>Xylella fastidiosa</i>), Bacterial Blast (<i>Pseudomonas syringae</i>), Bacterial Canker (<i>Pseudomonas</i> spp), Bacterial Leaf Blight (<i>Xanthomonas axonopodis</i> pv. Allii), Bacterial Speck (<i>Pseudomonas syringae</i>), Bacterial Spot (<i>Xanthomonas</i> spp), Bacterial Wilt (<i>Ralstonia solanacearum</i>), Black leg /bacterial soft rot/Aerial Stem Rot (<i>Erwinia carotovora</i>), Black Rot (<i>Xanthomonas campestris</i>), Brown Spot (<i>Pseudomonas syringae</i>), Citrus Canker (<i>Xanthomonas citri</i>), Common Blight (<i>Xanthomonas phaseoli</i>), Fire blight (<i>Erwinia amylovora</i>), Garlic Blotch (<i>Pseudomonas marginalis</i>), Hazelnut Bacterial Blight (<i>Xanthomonas arboricola</i> pv. <i>Corylina</i>), Onion Soft Rot (<i>Burkholderia gladioli</i>), Walnut Blight (<i>Xanthomonas arboricola</i> pv. <i>Juglandis</i>)	24-45	50 – 200 by ground 7-20 by air	Apply in sufficient water to provide full coverage Begin applications prior to, or at the first signs of disease Continue applications on a 7-day interval while environmental conditions favor disease development

As with any crop-protection product, always read and follow label directions.

Sym-Agro, Inc
 111 South Court Street, Ste 206
 Visalia, CA 93291
www.sym-agro.com